

Article

Asian-Australasian Journal of Animal Sciences (AJAS) 2014; 27(9): 1293-1302.

<https://doi.org/10.5713/ajas.2013.13766>

Energy Requirements for Maintenance and Growth of Male Saanen Goat Kids

A. N. Medeiros¹, K. T. Resende¹, I. A. M. A. Teixeira¹, M. J. Araújo^{2, *}, E. A. Yáñez³, A. C. D. Ferreira⁴**Affiliations** ▶

*Corresponding Author: Marcos Jácome de Araújo.

Tel: +55-89-35622711, Fax: +55-89-35621866, E-mail: marcos_jacome@yahoo.com.br

Department of Animal Science, Universidade Federal da Paraíba (UFPB), Areia, PB 58397-000, Brazil

Received November 27, 2013 Revised January 21, 2014 Accepted April 28, 2014

Copyright © 2014 by Asian-Australasian Journal of Animal Sciences

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License <http://creativecommons.org/licenses/by-nc/3.0/> which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.**Abstract**Go to :

The aim of study was to determine the energy requirements for maintenance and growth of forty-one Saanen, intact male kids with initial body weight (BW) of 5.12±0.19 kg. The baseline (BL) group consisted of eight kids averaging 5.46±0.18 kg BW. An intermediate group consisted of six kids, fed for *ad libitum* intake, that were slaughtered when they reached an average BW of 12.9±0.29 kg. The remaining kids (n = 27) were randomly allocated into nine slaughter groups (blocks) of three animals distributed among three amounts of dry matter intake (DMI; *ad libitum* and restricted to 70% or 40% of *ad libitum* intake). Animals in a group were slaughtered when the *ad libitum*-treatment kid in the group reached 20 kg BW. In a digestibility trial, 21 kids (same animals of the comparative slaughter) were housed in metabolic cages and used in a completely randomized design to evaluate the energetic value of the diet at different feed intake levels. The net energy for maintenance (NE_m) was 417 kJ/kg^{0.75} of empty BW (EBW)/d, while the metabolizable energy for maintenance (ME_m) was 657 kJ/kg^{0.75} of EBW/d. The efficiency of ME use for NE maintenance (k_m) was 0.64. Body fat content varied from 59.91 to 92.02 g/kg of EBW while body energy content varied from 6.37 to 7.76 MJ/kg of EBW, respectively, for 5 and 20 kg of EBW. The net energy for growth (NE_g) ranged from 7.4 to 9.0 MJ/kg of empty weight gain by day at 5 and 20 kg BW, respectively. This study indicated that the energy requirements in goats were lower than previously published requirements for growing dairy goats.

Keywords: Body Composition; Comparative Slaughter; Dairy Goats; Feed restriction; Heat Production**INTRODUCTION**Go to :

Understanding energy requirements for goats is important to the rational design of animal production systems so as to provide balanced diets that precisely meet the needs of the animals. Most studies on the nutrition of goats have been published quite recently. A still widely used publication on nutrient requirements of goats was published by the US National Research Council (NRC, 1981). Nevertheless, the diets based on the recommendations of this committee do not always accurately estimate the feed intake and performance. Causes for this low prediction might be due to these generalities have been established primarily with cattle and sheep. Several years after the publication of the NRC (1981) system, recommendations more specific for goats were developed and published for other feeding systems (AFRC, 1998; Sahlu et al., 2004; INRA, 2007; NRC, 2007).

The system of Institute for Goat Research-Langston University (IGR system) was almost integrally adopted by the recently published NRC (2007) system for goats. In this system metabolizable energy (ME) requirement were determined by regressing estimates of intake against levels of production. An attribute of this approach is relatively large numbers of observations available for use, but there are disadvantages as well (Sahlu et al., 2009). Both NRC (2007) and Sahlu et al. (2004) indicated areas in which additional research is needed to more accurately describe energy requirements of goats for improved feeding management practices. Energy requirements of NRC (2007) are for ME. To

eventually advance to a net energy system, [Sahlu et al. \(2009\)](#) suggested that research is needed to address body composition and efficiency of ME utilization for different physiological functions.

More importantly, methodological differences could be an important source of variance of requirements estimates. Thus, for energy requirements of goats to be eventually established, a greater number of detailed studies of energy requirements and efficiencies of utilization are required. In addition, correct estimation of their energy requirements in each unique condition is necessary to optimize diet composition or to determine optimum stocking rates of goats. Therefore, this study was carried out to determine the energy requirements for maintenance and growth of Saanen goat kids using the comparative slaughter technique.

MATERIAL AND METHODS

Go to : 

Study location

This study was conducted at the Goat Center of Universidade Estadual Paulista (UNESP), in Jaboticabal, São Paulo, Brazil (21°14'05" S and 48°17'09" W). Mean minimum and maximum temperatures observed during the experimental period were 16.3°C and 30.3°C, respectively. Humane animal care and handling procedures were followed according to the university's animal care committee.

Animals, dietary treatments and feeding plan

Intact, male Saanen kids (n = 41) with average initial body weight (BW) of 5.12±0.19 kg and an average age of 16 days were used in this study. After birth, the kids were identified and housed in individual pens (0.50 m×1.00 m) protected from rain and wind with free access to water. During the first three days, the kids received only colostrum. From the 4th to 40th day of life, cow milk was provided at a maximum quantity of 1.5 liter, twice a day, at 0700 and 1700 h. From the 41st to the 48th day of life, the kids were suckled only in the afternoon, receiving 750 mL of milk. The milk intake of animals fed *ad libitum* was used to determine the intake of 40% and 70% of *ad libitum* intake. After 7 days of age, all kids began receiving the experimental diet for *ad libitum* intake and they were offered enough feed to ensure that there was about 20% feed refusals each day. The experimental diet ([Table 1](#)) was formulated according to [AFRC \(1998\)](#) recommendations. For all calculations and measurements, the dry matter intake (DMI) from milk during the suckling period and the DMI from solid feed during throughout the experiment were considered to totalize the DMI. During the trials, animals were fed twice a day, at 0700 and 1700 h.


Table 1

Ingredient and chemical composition of the diet

Item	Value
Dietary ingredient (g/kg dry matter)	
Dehydrated corn plant ¹	468.8
Cracked corn grain	259.1
Soybean meal	193.2
Molasses	42.9
Soybean oil	8.1
Mineral supplement ²	19.9
Limestone	8.0
Chemical composition	
Dry matter	908.9
Crude protein	164.6
Ether extract	13.7
Neutral detergent fiber	475.0
Acid detergent fiber	104.0
Lignin	33.7
Gross energy (MJ/kg of dry matter)	17.45

¹ Whole corn plants (60% to 70% moisture) chopped when the kernel milk line was approximately two-thirds of the way down the kernel. Chopped material was air dried for approximately 72 h or until it reached 8% to 10% moisture and then ground to pass through a 4-mm screen.

² Composition, per kg, as fed-basis: 190 g of Ca; 73 g of P; 62 g of Na; 90 g of Cl; 44 g of Mg; 30 g of S; 1,350 g of Zn; 340 mg of Cu; 940 mg of Mn; 1,064 mg of Fe; 3 mg of Co; 16 mg of I; 18 mg of Se; 730 mg of F (maximum).

 [Download Table](#)

Twenty-seven kids were randomly allocated to three amounts of DMI: *ad libitum* and intake of 70% or 40% of *ad libitum* intake. Kids were pair fed in nine slaughter groups (blocks). A slaughter group consisted of one kid from each treatment and was slaughtered when the *ad libitum* treatment kid reached 20 kg BW. The 40% and 70% of *ad libitum* intake amounts were determined daily based on DMI of the kids in the *ad libitum*-group on the previous day. Six kids were assigned to intermediate slaughter group that were fed *ad libitum* and slaughtered when they reached 12.9 ± 0.29 kg BW.

Chemical analyses

The feed ingredients were dried at 60°C to 65°C for 72 h and ground through a 1-mm screen using a Wiley mill (Arthur H. Thomas Co., Philadelphia, PA, USA). They were analyzed to determine the content of DM (AOAC, 1990; method number 930.15), fat (based on weight loss of the dry sample upon extraction with petroleum ether in a Soxhlet extraction apparatus for 6 h; AOAC, 1990, method number 930.15), crude protein (CP) (Nitrogen analysis performed via Dumas combustion using LECO FP-528LC, LECO Corp., St. Joseph, MI, USA; Etheridge et al., 1998), ash (complete combustion in a muffle furnace at 600°C for 6 h; AOAC, 1990, method number 924.05), neutral detergent fiber with amylase and without sulfite (Van Soest et al., 1991), acid detergent fiber (Goering and Van Soest, 1970), and gross energy (GE) using a bomb calorimeter (Parr Instrument Co., Moline, IL, USA).

Sample collection and digestibility trial

A digestibility trial with a completely randomized design was conducted using 21 kids (mean BW at start: 19.4 ± 3.9 kg; same animals of the comparative slaughter trial) at three levels of intake. This trial was conducted when the animals were only receiving solid feed. This was chosen due to carrying out a digestibility assay with very young animals is complicated because of the small amount and consistency of the feces, and the level of stress to which young animals would be subjected. Thus, the daily milk and solid diet intakes were recorded separately and analyzed (diet offered andorts) for DM, CP, and GE. A general relationship between ME and GE for milk ($ME_{\text{milk}} = 0.93 \times GE_{\text{milk}}$; NRC, 2001) was obtained from the literature, and using this factor the ME intake (MEI) from milk could be calculated. The MEI from solid could be calculated using the data from digestibility trial. Then the total MEI for the whole experiment was obtained as the sum of daily MEI from milk and daily MEI from solid diet.

The animals were housed in individual metabolic cages when they were at approximately 70 d of age. Their feed intake and feed refusals were recorded, and their feces and urine were collected for 7-d after a 5-d adaptation period to each amount of intake. Urine was acidified daily with 20 mL of 6 M HCl. Feeds and feed refusals were sampled daily, and the samples were stored at -20°C. Feces and urine were weighed daily, and a 10% total output was sampled and stored at -20°C. Composites of the feed, feed refusals, and feces were dried at 60°C to 65°C for 72 h and ground through a 1-mm screen using a Wiley mill. Composites of urine were passed through a sieve to remove the large particles, and a subsample was taken for N determination. The GE was determined for feeds, feed refusals, feces, and urine using a bomb calorimeter (Parr Instrument Co., USA). To measure urine energy, samples were oven dried for two days at 55°C in polyethylene capsule prior to combustion. The known heat of combustion per gram of capsule material was subtracted from the total heat observed to obtain the sample energy content. Digestible energy (DE) was computed from the GE of the feeds, feed refusals, and feces. The DE values were regressed on daily DMI (g/d) of the goats in the digestibility trial, and the regression was subsequently used in the comparative slaughter trial to estimate the DE content and intake. Dietary ME was calculated using the GE intake minus the fecal

and urinary losses and losses by gaseous products of digestion (Blaxter and Clapperton, 1965), as follows in the Eq. 1:

(1)

$$\text{ME (MJ/kg of DM)} = [\text{GEI} - (\text{GE}_f + \text{GE}_u + \text{EGPD})]$$

In which ME = metabolizable energy; GEI = GE intake; GE_f = GE of feces; GE_u = GE of urine; and EGPD = energy of gaseous products of digestion ($\text{GPD} \times \text{GEI} / 100$); $\text{GPD} = [4.28 + (0.059 \times \text{GE digestibility})]$.

Slaughtering and body composition determination

Before slaughter, shrunk body weight (SBW) was measured as the BW after 16 h without feed and water. At slaughter, kids were stunned with an electric shock and killed by exsanguinations using conventional humane procedures. Their blood was weighed and sampled. The body was separated into individual components, which were then weighed separately. The components included internal organs (liver, heart, lungs and trachea, tongue, kidneys, and spleen), emptied and cleaned digestive tract (rumen, reticulum, omasum, abomasum, and small and large intestines), hide, head, feet, and carcass. The digestive tract was weighed before and after emptying and flushing with water to determine the empty BW (EBW), which was calculated as the SBW at slaughter minus digestive tract contents.

All body components were initially frozen at -6°C and then cut into small pieces, ground with a large screw grinder through a plate with 0.32-cm holes, and mixed via 2 additional passes through the grinder. After grinding and homogenization, the samples were collected, frozen again, and freeze dried for DM determination. These samples, of approximately 30 g, were analyzed for fat, CP, and GE as described previously.

Calculation of the initial body composition and retention of nutrients

Initial empty body composition was estimated from the average composition of the initial slaughter group, called the baseline (BL) group, which was composed of eight kids averaging 5.46 ± 0.18 kg BW and 15 d old. A regression equation was developed from BL animals to determine the initial EBW from the BW (BW; Eq. 2). The empty body fat (EBF; Eq. 3), empty body protein (EBP; Eq. 4) and empty body energy (EBE; Eq. 5) were estimated from the EBW for each animal the average body composition of BL animals.

(2)

$$\text{EBW (kg)} = 2.96(\pm 0.36) + 0.38(\pm 0.07) \times \text{BW (kg)}$$

$$(\text{R}^2 = 0.84; \text{RMSE} = 0.03)$$

(3)

$$\text{Log}_{10} \text{EBF (g)} = -4.18(\pm 0.85) + 9.49(\pm 1.22) \times \text{log}_{10} \text{EBW (kg)}$$

$$(\text{R}^2 = 0.91; \text{RMSE} = 0.02)$$

(4)

$$\text{Log}_{10} \text{EBP (g)} = 1.91(\pm 0.50) + 1.46(\pm 0.72) \times \text{log}_{10} \text{EBW (kg)}$$

$$(\text{R}^2 = 0.41; \text{RMSE} = 0.01)$$

(5)

$$\text{Log}_{10} \text{EBE (MJ/kg EBW)} = -2.62(\pm 0.89) + 5.88(\pm 1.26) \times \text{log}_{10} \text{EBW (kg)}$$

$$(\text{R}^2 = 0.78; \text{RMSE} = 0.02)$$

Rates of EBW gain (EWG, kg/d) and of body fat, protein and energy gain were calculated as the difference between the initial and final weights of the respective body components, divided by number of days in the trial. The caloric values of retained fat and protein were assumed to be 39.3 and 23.6 MJ/kg (ARC, 1980), respectively.

Energy requirements for maintenance and growth

Heat production (HP, kJ/kg^{0.75} of EBW) was calculated as the difference between MEI (average MEI from milk during the suckling period+average MEI from solid feed; MEI, kJ/kg^{0.75} of EBW) and retained energy (RE) (kJ/kg^{0.75} of EBW). The antilog of the intercept of the linear regression between the log of HP and MEI was used to estimate the maintenance requirement for net energy (NE_m; kJ/kg^{0.75} of EBW; Lofgreen and Garrett, 1968). The maintenance requirement for ME (ME_m, kJ/kg^{0.75} of EBW) was computed by iteratively solving the semilog linear regression equation until HP was equal to MEI. Linear regressions of the log of HP on MEI were used to calculate the energy utilization for maintenance (k_m), which was computed as NE_m/ME_m. The slope of the regression of RE on MEI was assumed to be the partial efficiency of energy for growth (k_g).

The NE requirements for growth (NE_g) were calculated using only animals fed for *ad libitum* intake: eight kids slaughtered at 5.46±0.18 kg BW (BL), six kids slaughtered at 12.9±0.29 kg BW (intermediate slaughter), and nine kids slaughtered at 21.03±0.36 kg BW. Estimation of the composition for gain was obtained in two phases. In the first phase, logarithms of allometric equations (Eq. 6) were calculated to predict the protein, fat, or energy concentration from EBW (Table 5) (ARC, 1980):


Table 5

Logarithmic allometric equations to estimate the body composition of Saanen intact, male kids (n = 23)

Logarithmic allometric equation	R ²	RMSE	p-value	Body weight (kg) ¹			
				5	10	15	20
EBW (kg) = 0.86(±0.17)+0.80(±0.01)×SBW (kg)	0.99	0.37	<0.0001	4.85	8.84	12.84	16.83
Log ₁₀ fat (g/kg of EBW) = 1.54(±0.08)+1.34(±0.08)×log ₁₀ EBW (kg)	0.92	0.09	<0.0001	59.91	73.70	83.81	92.02
Log ₁₀ protein (g/kg of EBW) = 2.20(±0.01)+1.03(±0.01)×log ₁₀ EBW (kg)	0.99	0.01	<0.0001	169.79	173.65	176.08	177.88
Log ₁₀ energy (MJ/kg of EBW) = 0.69(±0.04)+1.16(±0.04)×log ₁₀ EBW (kg)	0.98	0.04	<0.0001	6.37	7.01	7.43	7.76

RMSE, root mean square error; EBW, empty BW.

¹ Values were calculated from the equations.

 Download Table

(6)

$$\text{Log}_{10} (\text{component amount}) = a + [b \times \text{log}_{10} (\text{EBW, kg})]$$

In which the component amount is the nutrient amount in the EBW (Table 5). In the second phase, equation (Eq. 6) was differentiated based on the EBW to compute estimates of the composition of the gain at various EBW (Eq. 7):

(7)

$$\text{Component} = b \times 10^a \times \text{EBW}^{(b-1)}$$

In which component is the amount of nutrient or energy concentration per unit of EBW gain (g/kg of gain or MJ/kg of gain) and a and b are parameters determined from a linear regression (Eq. 6). The NE_g requirements of the gain in BW were obtained through the conversion of EBW into BW. The weight gain composition values were divided by the correction factor generated from the relationship between BW and EBW.

Statistical analyses

The data of maintenance requirements were analyzed as a randomized block design using a mixed model with the fixed effects of amounts of intake ($df = 2$), random effect of slaughter groups ($df = 8$) and random residual error using PROC MIXED (SAS Inst. Inc., Cary, NC, 9.0). When significant, means across amounts of intake were compared using protected fisher's least significant difference test (i.e., the DIFF option of the LSMEANS statement). Data of requirements of growth were analyzed as a completely randomized design and linear regressions analyses using PROC REG of SAS. Residuals were plotted against the predicted values to check the model assumptions regarding the homoscedasticity, independence, and normality of the errors. A data point was deemed to be an outlier and removed from the database if the Studentized residual was outside the ± 2.5 range values.

RESULTS

Go to :

Performance and body composition

Intake influenced body composition (% EBW) of the animals (Table 2). The fat content (% EBW) for *ad libitum* feed-intake animals was greater than those of the 70% and 40% feed-intake animals. Fat deposited in the gain was lower ($p < 0.01$) for 40% feed-intake group compared those in the *ad libitum* and 70% feed-intake groups. Protein and protein in gain were greater for animals in the 40% feed-intake group than those in the *ad libitum* and 70% feed-intake group. The water content for 70% and 40% feed-intake was greater than that of the *ad libitum* feed-intake group. There was no difference in the ash concentration (% EBW) between animals submitted to the different amount of feed intake ($p > 0.05$). Animals in the *ad libitum* group had more energy deposited in the empty body ($p < 0.001$) compared with those fed the 70% and 40% amounts.

Table 2

Performance and body composition of intact, male Saanen kids at baseline (BL) and intermediate slaughter (IS) and subjected to three amounts of feed intake


Variable	BL	IS	Amount of feed intake ¹			SEM	p-value
			<i>Ad libitum</i> %	70%	40%		
Performance							
Number of animas	8	6	9	9	9	-	-
Days on feed	-	51	76	76	76	-	-
Days suckling	-	-	34	34	34	-	-
Final age, days	15	66	91	91	91	-	-
Initial BW (kg)	5.46	5.07	5.04	5.01	5.03	-	-
Final BW (kg)	5.46	12.90	21.03a	16.18b	11.07c	0.42	<0.0001
SBW (kg)	5.21	12.25	19.84a	15.57b	10.61c	0.31	<0.0001
EBW (kg)	5.03	10.61	16.71a	12.94b	8.69c	0.26	<0.0001
ADG (g/d)	-	153.60	213.44a	148.33b	79.92c	6.67	<0.0001
EWG (g/d)	-	112.28	157.33a	107.46b	50.75c	4.48	<0.0001

Variable	BL	IS	Amount of feed intake ¹			SEM	p-value
			<i>Ad libitum</i> %	70%	40%		
Body composition (% EBW)							
Fat	5.99	8.91	8.92 ^a	7.02 ^b	4.84 ^c	0.54	0.0003
Protein	16.96	17.65	17.72 ^b	17.93 ^b	18.54 ^a	0.16	<0.006
Water	72.42	68.71	68.41 ^b	70.22 ^a	71.69 ^a	0.56	0.003
Ash	4.63	4.72	4.95	4.83	4.92	0.08	0.53
Fat in gain	-	12.45	10.71 ^a	8.58 ^a	5.11 ^b	0.95	0.002
Protein in gain	-	18.46	18.14 ^b	18.68 ^b	20.91 ^a	0.26	<0.001
Energy (MJ kg/EBW)	6.38	7.50	7.66 ^a	6.89 ^b	6.13 ^c	0.22	0.0006

SEM, standard error of the mean; BW, body weight; SBW, shrunk body weight; EBW, empty body weight; ADG, average daily gain; EWG, empty weight gain.

¹ AL and restricted to 70% or 40% of the *ad libitum* intake. Animals in each group were slaughtered when the kid in the *ad libitum* treatment reached 20 kg. Values for IS and BL are for reference only; they were not used in the statistical.

^{a,b,c} Means within a row with different superscripts are significantly different ($p < 0.05$).

 Download Table

Digestibility trial

It was observed that the MEI (MJ/d) and DMI (g/d) decreased as the feed restriction (70% and 40% feed-intake groups) increased ($p < 0.0001$; Table 3). The values of DE and GE digestibility were greater for animals in the 40% feed-intake group than those in the *ad libitum* and 70% feed-intake groups which did not differ. The metabolizability and efficiency of DE used for ME of the diet showed no difference ($p > 0.05$) among amount of feed intake.

Table 3

Dry matter intake (DMI), metabolizable energy intake (MEI) and energy partitioning of Saanen kids subjected to three amounts of feed intake in a digestibility trial


Variable	Amount of feed intake ¹			SEM	p-value
	<i>Ad libitum</i> %	70%	40%		
Number of animals	7	7	7	-	-
Body weight (BW, kg) ²	19.40 ^a	14.78 ^b	10.06 ^c	0.32	<0.0001
Empty body weight (EBW, kg) ³	10.79 ^a	9.00 ^b	6.76 ^c	0.15	<0.0001
DMI (g/d)	943.72 ^a	621.16 ^b	368.78 ^c	40.03	<0.0001
DMI (g/kg ^{0.75} of EBW)	163.67 ^a	122.29 ^b	88.40 ^c	7.97	<0.0001
MEI (MJ/d)	9.74 ^a	6.55 ^b	4.08 ^c	0.57	<0.0001
MEI (MJ/kg ^{0.75} of EBW)	1.69 ^a	1.28 ^b	0.97 ^b	0.06	<0.0001
Gross energy digestibility (%)	67.71 ^b	69.52 ^b	73.50 ^a	1.45	0.01
Digestible energy (DE, MJ/kg of DM)	11.81 ^b	12.13 ^b	12.82 ^a	0.25	0.01
Metabolizable energy (ME, MJ/kg of DM)	10.31	10.52	10.98	0.29	0.16
Metabolizability (q _m)	59.06	60.28	62.95	1.67	0.16
ME/DE	87.23	86.68	85.52	0.93	0.32

SEM, standard error of the mean.

¹ *Ad libitum* and restricted to 70% or 40% of the *ad libitum* intake.

^{2,3} Average BW and EBW of animals during metabolism trial.

^{a,b,c} Means within a row with different superscripts are significantly different ($p < 0.05$).

 Download Table

The resulting regression equations (Eq. 8 and 9) were used in the comparative slaughtering experiment to estimate the DE (MJ/kg of DM) and ME (MJ/kg of DM) content and then the MEI for the entire experimental period. Intake level affected total DMI and total MEI ($p < 0.05$) and the energy balance of animals throughout the experiment (Table 4). Following the same trend seen in the digestibility trial (Table 3), reductions in the total DMI (g/d) and MEI (MJ/d) were significantly different ($p < 0.001$) among the intake levels.

Table 4

Dry matter intake (DMI), metabolizable energy intake (MEI) and energy balance of Saanen intact, male kids subjected to three amounts of feed intake throughout the experimental period

Variable	Amount of feed intake ¹			SEM	p-value
	<i>Ad libitum</i> %	70%	40%		
Body weight (BW, kg) ²	13.04a	10.59b	8.05c	0.15	<0.0001
Empty body weight (EBW, kg) ³	10.80a	8.90b	6.78c	0.13	<0.0001
DMI (g/d)					
As solid feed	435.43a	290.14b	185.62c	18.55	<0.0001
As milk	162.07a	119.18b	69.31c	3.84	<0.0001
Total	597.50a	409.32b	254.93c	17.66	<0.0001
MEI (MJ/d)					
As solid feed	4.66a	3.13b	2.02c	0.19	<0.0001
As milk	4.20a	3.09b	1.80c	0.09	<0.0001
Total	8.86a	6.22b	3.82c	0.18	<0.0001
MEI (MJ/kg ^{0.75} of EBW)					
As solid feed	0.80a	0.62b	0.48c	0.03	<0.0001
As milk	0.72a	0.61b	0.43c	0.02	<0.0001
Total	1.53a	1.23b	0.91c	0.05	<0.0001
Retained energy (MJ/kg ^{0.75} of EBW)	0.23a	0.16b	0.08c	0.01	<0.0001
Heat production (MJ/kg ^{0.75} of EBW)	1.30a	1.06b	0.82c	0.04	<0.0001

SEM, standard error of the mean.

¹ *Ad libitum* and restricted to 70% or 40% of the *ad libitum* intake.

^{2,3} Average BW and EBW of animals during throughout study.

a,b,c Means within a row with different superscripts are significantly different (p<0.05).

[Download Table](#)

(8)

$$DE \text{ (MJ/kg DM)} = 13.08(\pm 0.42) - 0.001(\pm 0.0006) \times DMI \text{ (g/d)}$$

(9)

$$ME \text{ (MJ/kg DM)} = 11.04(\pm 0.49) - 0.0007(\pm 0.0006) \times DMI \text{ (g/d)}$$

Retained energy

Animals fed for *ad libitum* intake had greater RE (MJ/kg^{0.75} of EBW) and HP (MJ/kg^{0.75} of EBW) than those animals in the 70% and 40% feed-intake groups (p<0.001; Table 4). The relationship between the RE in the body of animals and RE predicted by the retention of protein and fat (RE_{p+f}) (Eq. 10) showed a high degree of precision and accuracy. This means that the RE in the body of animals can safely be estimated based on the retention of protein and fat. The percentage of RE deposited as protein (% RE_p) decreased exponentially as the content of RE in the gain (RE, MJ/kg of EWG) increased (Figure 1; Eq. 11, the data referring only to the restriction period).




Figure 1

Relationship between the percentage of retained energy deposited as protein (% RE_p) and the content of retained energy in the empty body weight gain (RE; MJ/kg of empty weight gain, EWG): % RE_p = 188.50(±19.74) × exp[0.1506(±0.01) × RE], R² = 0.85; p<0.001].

[Download Figure](#)

(10)

$$\text{RE (MJ/d)} = 0.005(\pm 0.006) + [1.00(\pm 0.007) \times \text{RE}_{p+f} \text{ (MJ/d)}]$$

$$(\text{R}^2 = 0.99)$$

(11)

$$\% \text{RE}_p = 188.50 (\pm 19.74) \times \exp [-0.1506 (\pm 0.01) \times \text{RE}]$$

$$(\text{R}^2 = 0.85)$$

Energy requirement for maintenance and growth

The relationship between HP and MEI is shown in Figure 2. In the present study, the NE_m was obtained through regression between the log of HP ($\text{kJ/kg}^{0.75}$ of EBW) and MEI ($\text{kJ/kg}^{0.75}$ of EBW) for growing Saanen goats (Eq. 12). The NE_m of Saanen kids (from 5 to 20 kg BW) was $417 \text{ kJ/kg}^{0.75}$ of EBW ($357 \text{ kJ/kg}^{0.75}$ of BW), while the ME_m that was calculated by iteration assuming HP to be equal to MEI at maintenance was $657 \text{ kJ/kg}^{0.75}$ of EBW ($562 \text{ kJ/kg}^{0.75}$ of BW). The partial efficiency of ME used for NE_m (k_m) found in this study through the NE_m/ME_m ratio was 0.64.




Figure 2

Relationship between the logarithm of heat production (HP) and the ME intake (MEI; $\text{kJ/kg}^{0.75}$ of empty BW) for Saanen intact male kids. $\text{Log}_{10} \text{HP} = 2.62(\pm 0.014) + 0.0003(\pm 0.00001) \times \text{MEI}$; $\text{R}^2 = 0.97$, $\text{RMSE} = 0.0152$, $p < 0.0001$. BW, body weight; ME, metabolizable energy; RMSE, root mean square error.

[Download Figure](#)

(12)

$$\text{Log}_{10} \text{HP (kJ/kg}^{0.75} \text{ of EBW)} = 2.62 \pm (0.014) + 0.0003(\pm 0.00001) \times \text{MEI kJ/kg}^{0.75} \text{ of EBW}$$

$$(\text{R}^2 = 0.97)$$

Considering the weight range of 5 to 20 kg BW, an increased of NE_g from 7.38 to 9.00 MJ/kg EWG was observed (Table 6). An increase of 35% in the fat content, 18% in the energy content, and 4.5% in the protein content were observed when the BW ranged from 5 to 20 kg (Table 6). The k_g value was calculated as the slope of the regression of RE ($\text{MJ/kg}^{0.75}$ of EBW) on MEI ($\text{MJ/kg}^{0.75}$ of EBW). The slope of the equation was different from zero ($p < 0.001$) and indicated a k_g value of 0.21. In the current study, using $k_g = 21\%$, the ME_g requirements ranged from 34.10 to 36.07 MJ/kg average daily gain (ADG) for Saanen goats with BW ranging from 5 to 20 kg, respectively.

Table 6

Estimates of fat, protein and energy concentrations in the equation to predict net requirements for gain of the empty body weight gain (EWG) at different body weight of Saanen intact male kids

Variable	Body weight (kg)				Equation of net requirements for gain ¹
	5	10	15	20	
EBW (kg)	4.85	8.84	12.84	16.83	
Fat (g/kg of EWG)	80.57	99.11	112.72	123.75	$46.74 \times \text{EBW}^{0.344}$
Protein (g/kg of EWG)	176.14	180.14	182.68	184.54	$166.03 \times \text{EBW}^{0.037}$
Energy (MJ/kg of EWG)	7.38	8.12	8.62	9.00	$5.74 \times \text{EBW}^{0.159}$

EBW, empty body weight.

¹ Component concentration = $b \times 10^a \times \text{EBW}^{(b-1)}$, in which a and b are constants determined from the equations in Table 5.

 Download Table

DISCUSSION

Go to : 

It has been reported that, as the animal's EBW increases and physiological maturity approaches, the deposition of muscle tissue is reduced and the deposition of fat is increased. This behavior is considered as the normal growth pattern of the animal, where fat tissue has later growth compared to the body growth (Lawrence and Fowler, 2002). The animals fed for *ad libitum* intake had an EBW approximately 23% higher than animals in the 70% feed-intake group. However, the body compositions were similar, indicating that changes in the proportions of tissues deposited in body also depend on the quality and utilization of feed that the animals were provided. The RE in the empty body of the animal is directly related to the amount of fat and protein deposited in the empty body (ARC, 1980). Therefore, it was expected that the amount of energy in the empty body of animals in the 70% feed-intake group would be similar to that of animals fed for *ad libitum* intake. However, due to the factors besides EBW and age, the animals that fed *ad libitum* had more energy deposited in the empty body. It is possible that animals in the 70% feed-intake group used more energy for the deposition of protein, since this variable did not differ between these animals and those fed *ad libitum*. A decrease in the water content of the body of animals was associated with an increase in the fat proportion, confirming an inverse relationship between these two body constituents. A non-significant difference in the ash concentration (% of EBW) between animals submitted to the different feed intake levels may be attributable to the fact that the ash and protein concentration changes a little with weight and/or age as related by Bezabih and Pfeffer (2003).

In general, the loss of energy through feces increased and DE decreased as feed intake increases due to the decrease in retention time of the digesta in the rumen and factors influencing the motility of the gastrointestinal tract (NRC, 2007). The efficiency of DE used for ME of the diet showed no difference among feed intake levels. However, this efficiency average was higher than that recommended by the NRC (1989), which considers a constant value of 82.0%. Using fixed values may under or overestimate the ME of the diet, depending on the intake level and amount of fiber in the diet.

The RE and HP values were higher in animals fed *ad libitum* than those in the 70% and 40% feed-intake group. Indicating that HP increased with increasing MEI and that the quality and quantity of feed can change the HP and ME_m requirements (Tolkamp, 2010). Goats are able to survive under severe feed restriction because their ability to minimize HP or energy expenditure when on a low nutritional plane such as near maintenance and fasting. It has been suggested that goats may be more capable of decreasing energy expenditure in such conditions compared with other ruminant species (Silanikove, 2000).

The RE_p decreased exponentially as content of RE in the gain increased. This behavior was expected because the proportion of muscle and fat changes during the growth and development of the animal. Thus, the deposition of muscle tissue is greater at the beginning of the growth phase and decreases when the proportion of fat deposition increases (Colomer-Rocher et al., 1992). This agrees with the results reported in goats (Fernandes et al., 2007) and in cattle (Tedeschi et al., 2004; Chizzoti et al., 2007). This finding suggests that RE_p can be used to compute the partial efficiency of ME to NE for growth.

ME_m estimates (562 kJ/kg^{0.75} of BW) were lower than the average value of 580 kJ/kg^{0.75} of full BW for growing dairy goats and 624 kJ/kg^{0.75} of full BW for intact male goats reported by Sahu et al. (2004) and Luo et al. (2004e), respectively, which served as the basis for publication of the NRC

(2007). These differences might be partially attributable to experimental conditions. In the present study, the goats were housed in relatively small areas, such as metabolism chambers or crates, whereas goats in most of the publications from other studies were kept under more normal farm or production conditions, hence their greater activity. Another factor to consider in explaining lower ME_m estimates is that the kids were fed a combination diet (milk plus dry feed during the suckling period). According to NRC (1989), efficiencies of utilization of ME for maintenance and gain will be somewhat higher for milk than starter feeds; nevertheless, there are few data available on subject.

Our energy requirement estimates were different from recommendations from the older feeding system for goats (NRC, 1981; AFRC, 1998). The NRC (1981) recommends average values of 239 and 424 kJ/kg^{0.75} of BW for NE_m and ME_m , respectively. On the other hand, the AFRC (1998) suggests an average of 315 and 438 kJ/kg^{0.75} of BW for NE_m and ME_m , respectively. Possible factors responsible for this difference can be attributed to the different methods used to obtain maintenance energy, different breeds, experimental conditions and even the use of data from different species (cattle and sheep). In a study conducted by Fernandes et al. (2007) values of 323 and 494 kJ/kg^{0.75} of EBW for NE_m and ME_m , respectively were observed in studies with intact 3/4 Boer×1/4 Saanen goats (from 20 to 35 kg BW). Much of the variation found may be explained by differences in weight, maturity stage, and breed. Dairy animals have lower amounts of body fat and, therefore, different nutritional requirements.

Generally, the k_m value is calculated by the ARC (1980) from equation $k_m = 0.35 \times q_m + 0.503$, where q_m = metabolizability. Using this equation and the mean value of q_m measured in the present study (61%), the value of k_m would be 71.6%, which is approximately 12% greater than our k_m , which was 64%. Using the k_m obtained by ARC (1980) and the value of 417 kJ/kg^{0.75} of EBW as NE_m , the ME_m would be 582 kJ/kg^{0.75} of EBW, or approximately 12% less than the value found in this study (657 kJ/kg^{0.75} of EBW). Therefore, acceptance of k_m estimated by the ARC system would result in the underestimation of ME_m requirements. This underestimate has also been observed by other researchers with growing goats (Fernandes et al., 2007) and lambs (Galvani et al., 2008). The k_m found in this research is similar to the value suggested by the Cornell Net Carbohydrate and Protein System (CNCPS) for sheep (Cannas et al., 2004), as well as to that used by the Small Ruminant Nutrition System (SRNS) (Tedeschi et al., 2010) which assumed to be constant and equal to 64.4%.

The NE_g ranged from 7.38 to 9.00 MJ/kg EWG, considering the weight range of 5 to 20 kg BW. This increase in the NE_g may be attributable to the progressive increase in fat deposition with the increase in BW. For 3/4 Boer×1/4 Saanen animals with weights ranging from 20 to 35 kg of BW, Fernandes et al. (2007) observed values of 10.46 to 12.55 MJ/kg EWG. The NRC (1981) suggested 16.73 MJ/kg EWG for goats, regardless of BW. The requirements for goats of NRC (2007) were determined based on BW gain and did not consider the composition of tissue gain because insufficient data available in the literature. The animals studied by Fernandes et al. (2007) showed higher NE_g than animals in the current study. These differences in the NE_g requirements were probably due to the higher concentrations of fat in gain, and consequently, a higher concentration of energy in relation to protein and higher concentration of water in lean tissue than in adipose tissue (Luo et al., 2004e), due to the fact that animals were heavier and older.

In an attempt to isolate the influence of the difference in size between the digestive tracts of animals from different weights, a correction factor for each weight was used (BW/EBW), being 1.03, 1.13, 1.17, and 1.19, respectively for 5, 10, 15, and 20 kg BW. For animals with ADG of 100 g, the NE_g ranged from 0.71 to 0.76 MJ/d when the BW changed from 5 to 20 kg, while Fernandes et al. (2007) reported higher NE_g values (0.91 to 1.08 MJ/d). The AFRC (1998) recommended values from 0.66 to 1.15 MJ/d for animals with weights ranging from 5 to 20 kg of BW, respectively. These differences may be due to the changes that occurred in the composition of body gain, among other factors, as previously mentioned. In addition, another explanation is that the AFRC (1998) equations are not suited for all breeds, since they consider an absolute BW value, regardless the degree of maturity of

the animals. Recommendations for energy requirements of goats from [NRC \(2007\)](#) are based primarily on reports summarized by [Sahlu et al. \(2004\)](#) which did not consider the composition of tissue gain and the estimates were determined based on dietary requirement that is ME_g .

In the current study, the slope of the regression of RE on MEI was assumed to be the k_g which indicated a value of 0.21. This value is different to the value (0.48) estimated using the [AFRC \(1993\)](#) equation ($k_g = 0.006 + 0.78 \times q_m$) and to the value (0.38) estimated considering the equation cited by [Tedeschi et al. \(2010\)](#) ($k_g = 18.36 / [27 + 41 \times RE_p]$). According to these authors, the use of the combination of body composition and energy content of the diet might be a better approach in computing the k_m and k_g . Therefore, there is a need for new research into the energy requirements of goats, as well as to develop mathematical models that more efficiently represent the biological behavior of nutrient deposition.

CONCLUSION

Go to : 

In conclusion, our study indicated that NE_m was 417 kJ/kg^{0.75} of EBW, while the ME_m was 657 kJ/kg^{0.75} of EBW. The male Saanen goats have a requirement of NE_g ranging from 7.38 to 9.0 MJ/kg of EWG. The current study demonstrates that growing Saanen kids have nutritional requirements for energy lower than those that have been reported by most international feeding systems. Additional research is needed for other classes of goats fed under different ratios of concentrate to forage, and different management conditions and with a wider range of goat BW.

ACKNOWLEDGMENTS

Go to : 

We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brasília, DF, Brazil) and Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP, Brazil, Proc. 04/05833-1) for providing financial support.

REFERENCES


Go to : 



AFRC (Agricultural and Food Research Council). 1993. Energy and Protein Requirements of Ruminants. CAB International; Wallingford, UK:

AFRC (Agricultural and Food Research Council). 1998. The nutrition of goat. An advisory manual prepared by the AFRC technical committee on responses to nutrients. CAB International; Wallingford, UK:

ARC (Agricultural Research Council). 1980. The nutrient requirements of ruminant livestock. CAB International; Wallingford, UK:

AOAC (Association of Official Analytical Chemists). 1990. Official Methods of Analysis. 15th edition Arlington, VA, USA:

Bezabih M, Pfeffer E. 2003. Body composition and efficiency of energy and nutrient utilization by growing pre-ruminant Saanen goat kids. *Anim Sci* 77:155–163.
[Article](#) 

Blaxter KL, Clapperton JL. 1965. Prediction of the amount of methane produced by ruminants. *Br J Nutr* 19:511–522.
[Article](#)  [PubMed](#) 

Cannas A, Tedeschi LO, Fox DG, Pell AN, Van Soest PJ. 2004. A mechanistic model for predicting the nutrient requirements and feed biological values for sheep. *J Anim Sci* 82:149–169.

[Article](#) [PubMed](#)

- Chizzotti ML, Valadares Filho SC, Tedeschi LO, Chizzotti FHM, Carstens GE. 2007. Energy and protein requirements for growth and maintenance of F₁Nellore×Red Angus bulls, steers, and heifers. *J Anim Sci* 85:1971–1981.
[Article](#) [PubMed](#)
- Colomer-Rocher F, Kirton AH, Mercer GJK, Duganzich DM. 1992. Carcass composition of New Zealand Saanen goats slaughtered at different weights. *Small Rumin Res* 7:161–173.
[Article](#)
- Etheridge RD, Pesti GM, Foster EH. 1998. A comparison of nitrogen values obtained utilizing the Kjeldahl nitrogen and Dumas combustion methodologies (Leco CNS 2000) on samples typical of an animal nutrition analytical laboratory. *Anim Feed Sci Technol* 73:21–28.
[Article](#)
- Fernandes MHMR, Resende KT, Tedeschi LO, Fernandes JS Jr, Silva HM, Carstens GE, Berchielli TT, Teixeira IAMA, Akinaga L. 2007. Energy and protein requirements for maintenance and growth of Boer crossbred kids. *J Anim Sci* 85:1014–1023.
[Article](#) [PubMed](#)
- Galvani DB, Pires CC, Kozloski GV, Wommer TP. 2008. Energy requirements of Texel crossbred lambs. *J Anim Sci* 86:3480–3490.
[Article](#) [PubMed](#)
- Goering HK, Van Soest PJ. 1970. Forage Fiber Analyses (Apparatus, Reagents, Procedures, and Some Applications). *Agricultural Handbook*. 379ARS-USDA; Washington, DC, USA:
- INRA (Institut National de la Recherche Agronomique). 2007. Alimentation des bovins, ovins et caprins. Besoins des animaux. Valeurs des aliments. Editions Quae; Paris, France:
- Lawrence TLJ, Fowler VR. 2002. Growth of Farm Animals. 2th Edition CAB International; Wallingford, UK:
- Lofgreen GP, Garrett WN. 1968. A system for expressing net energy requirements and feed values for growing and finishing beef cattle. *J Anim Sci* 27:793–806.
[Article](#)
- Luo J, Goetsch AL, Sahlu T, Nsahlai IV, Johnson ZB, Moore JE, Galyean ML, Owens FN, Ferrell CL. 2004e. Prediction of metabolizable energy requirements for maintenance and gain of preweaning, growing and mature goats. *Small Rumin Res* 53:231–252.
[Article](#)
- NRC (National Research Council). 1981. Nutrient Requirements of Goats: Angora, Dairy and Meat Goats in Temperate and Tropical Countries. National Academy Press; Washington, DC, USA:
- NRC (National Research Council). 1989. Nutrient Requirements of Dairy Cattle. 6th revised Edition National Academy Press; Washington, DC, USA:
- NRC (National Research Council). 2001. Nutrient Requirements of Dairy Cattle. National Academy Press; Washington, DC, USA:
- NRC (National Research Council). 2007. Nutrient Requirements of Small Ruminants, Sheep, Goats, Cervids, and New World Camelids. The National Academic Press; Washington DC, USA:
- Sahlu T, Goetsch AL, Luo J, Nsahlai IV, Moore JE, Galyean ML, Owens FN, Ferrel CL, Johnson ZB. 2004. Nutrient requirements of goats: developed equations, other considerations and future

research to improve them. *Small Rumin Res* 53:191–219.

[Article](#) 

Sahlu T, Dawson LJ, Gipson TA, Hart SP, Merkel RC, Puchala R, Wang Z, Zeng S, Goetsch AL. 2009. ASAS Centennial Paper: Impact of animal science research on United States goat production and predictions for the future. *J Anim Sci* 87:400–418.

[Article](#)  [PubMed](#) 

Silanikove N. 2000. The physiological basis of adaptation in goats to harsh environments. *Small Rumin Res* 35:181–194.

[Article](#) 

Tedeschi LO, Fox DG, Guiroy PJ. 2004. A decision support system to improve individual cattle management. 1. A mechanistic, dynamic model for animal growth. *Agric Syst* 79:171–204.

[Article](#) 

Tedeschi LO, Cannas A, Fox DG. 2010. A nutrition mathematical model to account for dietary supply and requirements of energy and other nutrients for domesticated small ruminants: The development and evaluation of a Small Ruminant Nutrition System. *Small Rumin Res* 89:174–184.

[Article](#) 

Tolkamp BJ. 2010. Efficiency of energy utilization and voluntary feed intake in ruminants. *Animal* 7:1084–1092.

[Article](#) 

Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods of dietary fiber, neutral detergent fiber and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci* 74:3583–3597.

[Article](#)  [PubMed](#) 
