

Update on the distribution of *Diphylla ecaudata* Spix, 1823 (Mammalia, Chiroptera): New records from the Brazilian northeast

Patrício A. Rocha^{1,3*}, Mônica A. Pedroso², Anderson Feijó¹, Newton Gurgel Filho¹, Bruno A.T.P. Campos¹ and Stephen F. Ferrari^{1,3}

1 Universidade Federal da Paraíba, Departamento de Sistemática e Ecologia, Programa de Pós-Graduação em Ciências Biológicas (Zoologia), Cidade Universitária, CEP: 58051-900, Castelo Branco, João Pessoa, PB, Brazil.

2 Universidade de Mogi das Cruzes, Campus da Sede, Avenida Candido Xavier de Almeida e Souza 200, Vila Partenio, CEP: 08780-911, Mogi das Cruzes, SP, Brazil.

3 Universidade Federal de Sergipe, Departamento de Ecologia, Cidade Universitária Professor José Aloísio de Campos, Avenida Marechal Rondon, CEP: 49100-000, Jardim Rosa Elze, São Cristóvão, SE, Brazil.

* Corresponding author. E-mail: parocha2@yahoo.com.br

ABSTRACT: The subfamily Desmodontinae encompasses the only mammals known to feed exclusively on the blood of other vertebrates. This study provides records of the desmodontine bat *Diphylla ecaudata* from a major gap in its known distribution in northeastern Brazil. Specimens were captured at five localities, all associated with caves. Two of these sites are located in the state of Sergipe, two in Bahia, and one in Ceará. The record of *D. ecaudata* from Ceará is the first for this state.

DOI: 10.15560/10.6.1541

Phyllostomidae is one of the most diverse bat families in both morphological and ecological terms, covering the full diversity of chiropteran feeding niches, including species that are insectivorous, nectarivorous, frugivorous, carnivorous, omnivorous, and sanguivorous (Kunz 1982). The subfamily Desmodontinae is composed of the only three mammal species known to feed exclusively on the blood of other vertebrates— the Common Vampire Bat, *Desmodus rotundus* (É. Geoffroy, 1810), the White-winged Vampire Bat, *Diaemus youngii* (Jentink, 1893), and the Hairy-legged Vampire Bat, *Diphylla ecaudata* Spix, 1823 (Kwon and Gardner 2008).

Diphylla ecaudata occurs between México and southern Brazil, with records from Colombia, Venezuela, Ecuador, Peru, and Bolivia (Greenhall *et al.* 1984; Kwon and Gardner 2008). In Brazil, the species is found in all biomes (Paglia *et al.* 2012). There is an apparent lacuna between the central Amazon basin and southwestern South America, however, which suggests a potentially disjunct distribution for the species in this continent (Kwon and Gardner 2008). Prior to the present study, there was also an extensive gap in the known distribution of the species in the Brazilian Northeast. The present study provides records of *D. ecaudata* from five new localities (Figure 1), two in the Brazilian state of Sergipe, two in Bahia, and one in Ceará. In all five cases, the occurrence of *D. ecaudata* was associated with the presence of caves or mines.

The specimens were collected under the license 27524-1 (SISBIO) either in mist-nets, in open habitats, or by using hand nets, when captured in caves. All specimens were handled in accordance with the recommendations of Sikes *et al.* (2011). The voucher specimens were deposited in the mammal collection of the

Universidade Federal da Paraíba (UFPB), in João Pessoa (Paraíba, Brazil) and in Laboratory of Conservation Biology (LBC) of the Universidade Federal de Sergipe, in São Cristóvão (Sergipe, Brazil), after being fixed in 10% formaldehyde and preserved in 70% ethanol, with the subsequent extraction of the skull. A total of 13 specimens were collected between 2010 and 2012, five in two caves (Casa de Pedra and Miaba) in Sergipe (LBC 31, LBC 32, LBC 73, LBC 78, LBC 79), three in Paripiranga, Bahia (UFPB 6548, UFPB 6564, UFPB 6566), two in Boqueirão da Onça, Bahia (UFPB 6663, UFPB 6665), and three in Santa Quitéria, Ceará (UFPB 9255, UFPB 9256, UFPB 9257).

Casa de Pedra cave (10°50' S, 37°27' W) is located in the municipality of Itabaiana (Sergipe) and has a horizontal extension of around 200 m. The surrounding landscape is dominated by pastures and orchards, which are relatively dense along watercourses. In this cave, two individuals of *D. ecaudata* were spotted close (about 20 cm distance) of a group of approximately 200 individuals of *D. rotundus*. Miaba cave is located in the municipality of São Domingos (10°43' S, 37°37' W), close to the Vaza-Barris River, in an ecotone of the Atlantic Forest and Caatinga domains, known locally as the Agreste. A resident group of approximately 50 individuals of *D. ecaudata* was found at this site.

The Fim-do-Morro-do-Parafuso cave (10°41' S, 37°51' W) in Paripiranga, Bahia, is surrounded by grazing land and plantations of manioc and maize, with some fragments of shrubby caatinga mainly along the local rivers. Despite the large number of caves found in the area, 15 of which were inspected, *D. ecaudata* groups were found only in Fim-do-Morro-do-Parafuso. The second site in Ba-




FIGURE 1. Map showing locality records of *Diphylla ecaudata* in Central and South America (blue triangles represent the new records assigned here). For key to code numbers see Table 1.

hia, Boqueirão da Onça (09°52' S, 41°06' W), in the municipality of Sento Sé, is characterized by a complex of hills and valleys covered by an arboreal caatinga (deciduous trees of 10–12 m height) that forms a closed canopy in the wet season (December–April). In the valleys, where the individuals were captured, the rivers retain moisture, even during the dry season, which allows the formation of dense, evergreen vegetation.

In Ceará, *D. ecaudata* specimens were captured near an abandoned mine in Santa Quitéria (04°33' S, 39°46' W), a region dominated by open, shrubby caatinga on shallow, stony soils (IPECE 2009). The region encompasses a number of intermittent watercourses, which are associated with denser patches of semideciduous woodland.

Diphylla ecaudata is the smallest vampire bat, with a head and body length of 75–93 mm, forearm of 50–56 mm, and weight of 24–43 g (Greenhall et al. 1984). This species feeds preferentially on the blood of birds, with bites being observed on the margin of the cloaca, and on the tarsus and feet (Gardner 1977). It is distinguished from the other desmodontines by the presence of two lower molars and a much wider postorbital constriction. Externally, the uropatagium, legs, and feet of *D. ecaudata* are conspicuously hirsute, the thumb is short (< 13 mm), with no basal pad, the eyes are large, and the ears short and rounded (Allen 1896; Greenhall et al. 1983).

Diphylla ecaudata is found primarily in natural refuges,

in particular caves, where roosting sites may be selected according to environmental conditions (e.g., humidity, temperature) and availability of crevices (Greenhall et al. 1984; Altringham 1996). Almost half (39) of the 86 records of the species in the wild have been obtained from surveys of caves (Table 1). At both sites in Sergipe, *Diphylla* was found cohabiting with *D. rotundus*. This latter species is generally more common than *D. ecaudata* not only in the local caves (PAR, personal observation), but also in those from southern (Trajano 1985) and central Brazil (Bredt et al. 1999). The greater selectivity of roosts observed in *D. ecaudata* may, at least partly, account for this difference; *D. rotundus* can often be found in man-made structures (Gomes and Uieda 2004).

This study expands considerably the known distribution of *D. ecaudata* in the caatinga scrub of northeastern Brazil—previously restricted almost entirely to the state of Pernambuco and neighboring areas—and represents the first record of this species for the state of Ceará, where a total of 35 chiropteran species are now known to occur (Gurgel-Filho and Langguth, in press). These findings further support the conclusion of Kwon and Gardner (2008) that the highly disjunct distribution of the species (Figure 1) may simply be due to a lack of adequate inventories.

ACKNOWLEDGEMENTS: We are grateful to CAPES (PAR, AF, BATPC) for graduate fellowships and CNPq for a research grant to SFF (process no. 303994/2011-8) and PAR (501701/2013-3).

TABLE 1. Localities at which *Diphylla ecaudata* has been recorded. The code numbers refer to the points shown in Figure 1. Localities marked with an asterisk (*) are caves.

COUNTRY/ POINT	LAT-Y	LONG-X	STATE	LOCALITY	SOURCE
Mexico					
1	22.63333	-99.0333	Tamaulipas	Sierra Madre Oriental *	Elizalde-Arellano et al. (2007)
Venezuela					
2	10.45	63.96667	Sucre	21 km E of Cumaná	Handley (1976)
3	10.35	67.66667	Aragua	Portachuelo	Ojasti and Linares (1971)
4	8.716667	-71.45	Mérida	Cueva de la Azulita	Ojasti and Linares (1971)
Colombia					
5	11.283333	-74	Magdalena	Cacagualito	Allen (1900)
6	4.25	73.56667	Meta	Restrepo	Aellen (1970)
7	2.166667	-71.5	Vaupés	Cerro de las Pinturas	Wenzel et al. (1966)
Ecuador					
8	-1.466667	78.13333	Pastaza	Mera	Albuja (1983)
Peru					
9	-5.6	78.28333	Amazonas	10 Km by trail SE of La Peca	Graham and Barkley (1984)
10	-6.166667	77.26667	San Martín	Puca Tambo	Thomas (1926)
11	-8.283333	-74.65	Ucayali	Yarinacocha	Sanborn (1949)
12	-9.616667	74.93333	Huánuco	Panguana	Hutterer et al. (1995)
13	-10.35	-74.85	Pasco	Nevati	Tuttle (1970)
14	-11.783333	-72.7	Cusco	San Martín-3	Solari et al. (2001)
Brazil					
15	-7.45	73.68333	Acre	Parque Nacional da Serra do Divisor	Nogueira et al. (1999)
16	-8.766667	-63.9	Rondônia	Porto Velho (USNM 562686)	Mok et al. (1982)
17	2.05	-50.8	Amapá	Macapá*	Peracchi et al. (1984)
18	-2.533333	54.28333	Pará	Taperinha	Piccinini (1974)
19	-3.3	52.33333	Pará	Caverna Pedra da Cachoeira, Altamira *	Pinto-da-Rocha (1995)
20	-3.45	52.88333	Pará	Gruta Wukorangma, Medicilândia *	Pinto-da-Rocha (1995)
21	-3.45	52.88333	Pará	Gruta Upu Murem, Medicilândia *	Pinto-da-Rocha (1995)
22	6.016667	-50.3	Pará	Gruta do (Xis), Parauapebas *	Pinto-da-Rocha (1995)
23	7.716667	51.18333	Pará	Gradaús	Piccinini (1974)
24	-8.85	43.46667	Piauí	Parque Nacional da Serra das Confusões *	Gregorin et al. (2008)
25	-4.316667	-40.15	Ceará	Santa Quitéria*	This Study
26	-7.483333	36.28333	Paraíba	Fazenda Bravo, Cabaceiras	Feijó and Langguth (2011)
27	-7.166667	34.91667	Paraíba	Caverna da Onça, João Pessoa *	Feijó and Langguth (2011)
28	-7.983333	-36.05	Pernambuco	Pedra dos Pontais, Toritama	Astúa and Guerra (2008)
29	-8.266667	35.96667	Pernambuco	Parque Ecológico Municipal Professor João Vasconcelos, Caruaru	Sousa et al. (2004)
30	-8.333333	36.41667	Pernambuco	Sítio Coendou, Serra da Balança, Belo Jardim	Astúa and Guerra (2008)
31	-8.133333	36.36667	Pernambuco	Pedra do Caboclo, Barra do Farias, Brejo da Madre de Deus*	Astúa and Guerra (2008)
32	-8.266667	38.03333	Pernambuco	RPPN Maurício dantas, Betânia	Cruz et al. (2005)
33	-7.5	-39.7	Pernambuco	Exu	Mares et al. (1981)
34	-7.616667	39.78333	Pernambuco	Fazenda Maniçoba, Exu	Mares et al. (1981)
35	-7.6	39.83333	Pernambuco	Serra da Farinha, Sítio da Gritadeira, Exu	Astúa and Guerra (2008)
36	-9.633333	37.78333	Sergipe	Fazenda Mundo Novo, Canindé do São Francisco	Leal et al. (2013)
37	-10.68333	37.41667	Sergipe	Caverna Casa de Pedra, Itabaiana *	This Study
38	-10.78333	37.56667	Sergipe	Caverna da Miaba, São Domingos *	This Study
39	-10.66667	-37.85	Bahia	Caverna Morro do Parafuso, Paripiranga *	This Study
40	-8.966667	39.53333	Bahia	Rio São Francisco	Type localyte of <i>D. ecaudata</i>
41	-9.733333	41.86667	Bahia	Boqueirão da Onça, Sento Sé	This Study
42	-12.23333	41.61667	Bahia	Gruta da Fumaça, Iraquar*	Sbragia and Cardoso (2008)
43	-12.93333	-41.1	Bahia	Caverna Poço Encantado, Itaetê *	Gregorin and Mendes (1999)
44	-15.41667	39.33333	Bahia	Vale do Rio Pardo, Santa Luzia *	Faria et al. (2006)
45	-15.45	-39.65	Bahia	Vale do Rio Pardo, Pau Brasil *	Faria et al. (2006)
46	-12.35	44.96667	Bahia	Lapa do Manoel Lopes, São Desidério *	Pinto-da-Rocha (1995)
47	-14.38333	-46.3	Goiás	APA Nascentes do Rio Vermelho *	Esbérard et al. (2005)
48	-14.48333	-46.1	Goiás	Caverna Revolucionária, Mambaí *	Esbérard et al. (2001)
49	-15.53333	47.86667	Distrito Federal	Gruta da Saúva, Sobradinho*	Bredt et al. (1999)
50	-15.48333	-48.05	Goiás	Toca da Gameleira, Padre Bernardo*	Bredt et al. (1999)
51	-14.3833	-46.3	Goiás	Gruta Morro, Padre Bernardo*	Bredt et al. (1999)
52	-14.4833	-46.1	Distrito Federal	Gruta do Sal e Gruta Fenda II (Brazlândia, Fazenda Palestina) *	Bredt et al. (1999)
53	-18.9333	-48.3	Minas Gerais	Itacarambi	Trajano and Gimenez (1998)
54	-12.9333	-41.1	Minas Gerais	Uberlândia	Stutz et al. (2004)
55	-15.0833	-44.0833	Minas Gerais	Curvello	Almeida et al. (2002)

TABLE 1. Continued.

COUNTRY/ POINT	LAT-Y	LONG-X	STATE	LOCALITY	SOURCE
56	-19.8	44.18333	Minas Gerais	Fazenda Paraíso, Esmeraldas	Uieda (1992)
57	-20.55	-41.2333	Espírito Santo	Reserva Biológica dos Comboios, Linhares	Ruschi (1978)
58	-19.7	-39.9333	Espírito Santo	Reserva Biológica da Pedra Azul, Domingos Martins	Ruschi (1982)
59	-18.75	-44.4	Espírito Santo	Gruta do Limoeiro, Castelo *	Ruschi (1951)
60	-22.4833	-42.8833	Rio de Janeiro	Caverna Santana, Cantagalo *	Esbérard et al. (1998)
61	-22.9	-43.2333	Rio de Janeiro	Estação Ecológica do Paraíso	Esbérard (2007)
62	-21.9833	-42.3667	Rio de Janeiro	Reserva Biológica do Tinguá	Dias and Peracchi (2008)
63	-20.4	-41.0167	Rio de Janeiro	Rio de Janeiro	Uieda (1986)
64	-22.7833	-43.6833	Rio de Janeiro	Serra da Tiririca, Sítio Três Nascentes	Teixeira and Peracchi (1996)
65	-22.55	-43.4	Rio de Janeiro	Parque Estadual da Pedra Branca	Dias et al. (2002)
66	-22.9667	-43.4667	Rio de Janeiro	Universidade Federal Rural do Rio de Janeiro, Seropédica	Piccinini et al. (1991)
67	-22.8667	-43.3	São Paulo	Cotia	McNab (1969)
68	-23.6167	-46.9333	São Paulo	Iguapé	Pira (1904)
69	-24.55	-48.6667	São Paulo	Abismo do Tira Prosa, Ribeira *	Pinto-da-Rocha (1995)
70	-24.5833	-48.7	São Paulo	Caverna de Santana, Iporanga *	Pinto-da-Rocha (1995)
71	-24.5167	-48.7	São Paulo	Gruta Bethary de Baixo, Iporanga *	Pinto-da-Rocha (1995)
72	-24.55	-48.65	São Paulo	Ressurgência das Areias, Iporanga *	Pinto-da-Rocha (1995)
73	-24.7167	-47.55	São Paulo	Gruta do Alambari de Baixo, Iporanga *	Pinto-da-Rocha (1995)
74	-24.55	-48.6667	São Paulo	Gruta das Areias de Cima, Iporanga *	Pinto-da-Rocha (1995)
75	-24.5833	-48.7	São Paulo	Gruta das Areias de Baixo, Iporanga *	Pinto-da-Rocha (1995)
76	-24.4	-49	São Paulo	Vale do Ribeira, Iporanga *	Traiano (1985)
77	-25.0333	-49.0833	Paraná	Gruta São João, Adrianópolis *	Pinto-da-Rocha (1995)
78	-23.45	-51.25	Paraná	Gruta Fada, Parque Estadual de Campinhos *	Arnone and Passos (2007)
79	-25.4167	-50	Paraná	Gruta de Toquinhas, Rio Branco do Sul *	Pinto-da-Rocha (1995)
80	-24.5833	-48.5833	Paraná	Palmeira	Thomas (1900)
81	-25.15	-49.3	Paraná	Parque Estadual Mata do Godoy, Londrina	Reis et al. (2000; 2003; 2012)
82	-27.2167	-49.15	Santa Catarina	Blumenau	Cherem et al. (2004)
83	-24.75	-48.5333	Santa Catarina	Gruta de Botuvera, Botuvera *	Pinto-da-Rocha et al. (2001)
84	-27.6833	-48.7667	Santa Catarina	Vidal Ramos	Cherem et al. (2004)
85	-26.9167	-49.0833	Santa Catarina	Santo Amaro da Imperatriz	Cherem et al. (2004)

LITERATURE CITED

- Aellen, V. 1970. Catalogue raisonné des chiroptères de la Colombie. Muséum d'Histoire Naturelle de Genève. *Revue Suisse de Zoologie* 77(1): 1–37 (<http://biodiversitylibrary.org/page/40807622>).
- Albuja, V.L. 1983. *Murciélagos del Ecuador*. Escuela Politécnica Nacional, Departamento de Ciencias Biológicas. Quito. 285 pp.
- Allen, H. 1896. Notes on the vampire bat (*Diphylla ecaudata*) with special reference to its relationship with *Desmodus rufus*. *Proceedings of the United States National Museum* 18(1099): 769–777 (doi: 10.5479/si.00963801.18-1099.769).
- Allen, J.A. 1900. List of bats collected by Mr. H. H. Smith in the Santa Marta region of Colombia, with descriptions of new species. *Bulletin of the American Museum of Natural History* 13(2): 87–94.
- Almeida, E.O., E.C. Moreira, L.A.B. Naveda, G.P. Herrmann. 2002. Combate ao *Desmodus rotundus* (E. Geoffroy, 1810) na região cárstica de Cordisburgo e Curvelo, Minas Gerais. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 54(2): 117–126 (doi: 10.1590/S0102-09352002000200002).
- Altringham, J.D. 1996. *Bats, Biology and Behavior*. Oxford: University Press. 262 pp.
- Arnone, I.S. and F.C. Passos. 2007. Estrutura de comunidade da quiropterofauna (Mammalia, Chiroptera) do Parque Estadual de Campinhos, Paraná, Brasil. *Revista Brasileira de Zoologia* 24(3): 573–581 (doi: 10.1590/S0101-81752007000300008).
- Astúa, D. and D.Q. Guerra. 2008. Caatinga bats in the mammal collection of the Universidade Federal de Pernambuco. *Chiroptera Neotropical* 14(1): 326–338.
- Bredt, A., W. Uieda and E.D. Magalhães. 1999. Morcegos cavernícolas da região do Distrito Federal, centro-oeste do Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 16(3): 752–754 (doi: 10.1590/S0101-81751999000300012).
- Cherem, J.J., P.C. Simões-Lopes, S. Althoff and M.E. Graipel. 2004. Lista dos mamíferos do estado de Santa Catarina, sul do Brasil. *Mastozoologia Neotropical* 11(2): 151–184.
- Cruz, M.A.O.M., D.M. Borges, A. Langguth, M.A.N. Sousa, L.A.M. Silva, L.M.R.M. Leite, F.M.V. Prado, K.C.S. Verissimo and B.L.C. Moraes. 2005. Diversidade de mamíferos em áreas prioritárias para conservação da Caatinga; pp. 182–201, in: F.S. Araujo, M.J. Rodal and M.R.V. Barbosa (eds.). *Análise das Variações da Biodiversidade do Bioma Caatinga: Suporte a Estratégias Regionais de Conservação*. Brasília: Ministério do Meio Ambiente.
- Dias, D., A.L. Peracchi and S.S.P. Silva. 2002. Quirópteros do Parque Estadual da Pedra Branca, Rio de Janeiro, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 19(2): 113–140 (doi: 10.1590/S0101-81752002000600012).
- Dias, D. and A.L. Peracchi. 2008. Quirópteros da Reserva Biológica do Tinguá, estado do Rio de Janeiro, sudeste do Brasil (Mammalia: Chiroptera). *Revista Brasileira de Zoologia* 25(2): 333–369 (doi: 10.1590/S0101-81752008000200023).
- Elizalde-Arellano, C., J.C. López-Vidal, J. Arroyo-Cabrales, R.A. Medellín and J.W. Laundré. 2007. Food sharing behavior in the hairy-legged vampire bat *Diphylla ecaudata*. *Acta Chiropterologica* 9(1): 314–318 (10.3161/1733-5329(2007)9[314:FSBITH] 2.0.CO;2).
- Esbérard, C.E.L. 2007. Influência do ciclo lunar na captura de morcegos Phyllostomidae. *Iheringia, Série Zoologia* 97(1): 81–85 (doi: 10.1590/S0073-47212007000100012).
- Esbérard, C.E.L., L.F.S. Martins, R.C. Cruz, R.C. Costa, M.S. Nunes, E.M. Luz and A.S. Chagas. 1998. Aspectos da biologia de *Lonchorhina aurita* no estado do Rio de Janeiro (Mammalia, Chiroptera, Phyllostomidae). *Revista Bioikos* 11(1–2): 46–49.
- Esbérard, C.E.L., J.A.O. Motta, E.M. Calvo, V.M. Ferreira, J.C. Carvalho, C.C. Carvalho, C.R.P. Souza, E.A. Pires, G.M.V. Rosa, J.S. Reis, J.N. Araújo and K.E. Quege. 2001. Morcegos cavernícolas de Mambá e arredores, Goiás, Brasil. Anais do Speleo Brasil, 26th Brazilian Congress of Speleology.
- Esbérard, C.E.L., J.A.O. Motta and C. Perigo. 2005. Morcegos cavernícolas da Área de Proteção Ambiental (APA) Nascentes do Rio Vermelho, Goiás. *Revista Brasileira de Zootecias* 7(2): 311–325.
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from the Atlantic rainforest of Southern Bahia, Brazil. *Biota Neotropica* 6(2): 1–13 (doi: 10.1590/S1676-06032006000200022).
- Feijó, J.A. and A. Langguth. 2011. Lista de quirópteros da Paraíba, Brasil com 25 novos registros. *Chiroptera Neotropica* 17(2): 1055–1062.
- Gardner, A.L. 1977. Feeding habits; pp. 293–350, in: R.J. Baker, J.K. Jones and D.C. Carter (eds.). *Biology of bats of the New World family Phyllostomatidae. Part. II. Special Publications of Museum of Texas*

- Tech University. Lubbock: Texas Tech Press.
- Gomes, M. N. and W. Uieda. 2004. Abrigos diurnos, composição de colônias, dimorfismo sexual e reprodução do morcego hematófago *Desmodus rotundus* (E. Geoffroy) (Chiroptera, Phyllostomidae) no Estado de São Paulo, Brasil. *Revista Brasileira Zoologia* 21(3): 629–638 (doi: 10.1590/S0101-81752004000300025).
- Graham, G.L. and L.J. Barkley. 1984. Noteworthy records of bats from Peru. *Journal of Mammalogy* 65(4): 709–711.
- Greenhall, A.M., G. Joermann, U. Schmidt and M.R. Seidel. 1983. *Desmodus rotundus*. *Mammalian Species* 202: 1–6.
- Greenhall, A.M., U. Schmidt and G. Joermann. 1984. *Diphylla ecaudata*. *Mammalian Species* 227: 1–3.
- Gregorin, R. and L.F. Mendes. 1999. Sobre quirópteros (Emballonuridae, Phyllostomidae, Natalidae) de duas cavernas da Chapada Diamantina, Bahia, Brasil. *Iheringia, Série Zoologia* 86: 121–124.
- Gregorin, R., A.P. Carmignotto and A.R. Percequillo. 2008. Quirópteros do Parque Nacional da Serra das Confusões, Piauí, nordeste do Brasil. *Chiroptera Neotropical* 14(1): 366–383.
- Gurgel-Filho, N.M. & Langguth, A. In press. Pequenos mamíferos do Ceará (Marsupiais, Morcegos e Roedores Sigmodontíneos). *Revista Nordestina de Biologia* 12(2).
- Handley, C.O., Jr. 1976. Mammals of the Smithsonian Venezuelan Project. *Brigham Young University Science Bulletin, Biological Series* 20(5): 1–89 (<http://biodiversitylibrary.org/page/7798479>).
- Hutterer, R., N. Verhaagh, J. Diller and R. Poudlucky. 1995. An inventory of mammals observed at Panguana Biological Station, Amazonian Peru. *Ecotropica* 1(1): 3–20.
- IPECE. 2009. *Instituto de Pesquisa e Estratégia Econômica do Ceará, Perfil Básico Municipal: Santa Quitéria*. Accessible at http://www.ipece.ce.gov.br/publicacoes/perfil_basico/pbm-2013/Santa_Quiteria. Captured on 20 December 2013.
- Kunz, T.H. 1982. *Ecology of bats*. New York: Plenum Press. 779 pp.
- Kwon, M. and A.L. Gardner. 2008. Subfamily Desmodontinae; pp. 218–224, in: A.L. Gardner (ed.). *Mammals of South America, Volume 1, Marsupials, Xenarthrans, Shrews, and Bats*. Chicago: University of Chicago Press.
- Leal, E.S.B., S.M. Azevedo-Junior, F.V.P. Vila-Nova, D.Q. Guerra and W.R. Telino-Junior. 2013. Updated compilation of bat species (Chiroptera) for the Brazilian state of Sergipe, including new records. *Chiroptera Neotropical* 19: 1163–1178.
- Mares, M.A., M.R. Willig, K.E. Streilein and T.E. Lacher. 1981. The Mammals of northeastern Brazil: A preliminary assessment. *Annals of Carnegie Museum* 50(4): 81–137.
- McNab, B.K. 1969. The economics of temperature regulation in Neotropical bats. *Comparative Biochemistry and Physiology* 31(2): 227–268.
- Mok, W.Y., D.E. Wilson, L.A. Lacey, and R.C.C. Luizão. 1982. Lista atualizada de quirópteros da Amazônia brasileira. *Acta Amazonica* 12(4): 814–823.
- Nogueira, M.R., A. Pol and A.L. Peracchi. 1999. New records of bats from Brazil with a list of additional species for the chiropteran fauna of the state Acre, western Amazon basin. *Mammalia* 63(3): 363–368.
- Ojasti, J., and O.J. Linares. 1971. Adiciones a la fauna de murciélagos de Venezuela con notas sobre las especies del género *Didicurus* (Chiroptera). *Acta Biologica Venezuelica* 7(1–2): 421–441.
- Paglia, A.P., G.A.B. Fonseca, A.B. Rylands, G. Herrmann, L.M.S. Aguiar, A.G. Chiarello, Y.L.R. Leite, L.P. Costa, S. Siciliano, M.C.M. Kierulff, S.L. Mendes, V.C. Tavares, R.A. Mittermeier and J.L. Patton. 2012. Lista anotada dos mamíferos do Brasil, 2ª ed. *Occasional Papers in Conservation Biology* 6: 76 pp.
- Peracchi, A.L., S.D.L. Raimundo and A.M. Tannure. 1984. Quirópteros do Território Federal do Amapá, Brasil (Mammalia, Chiroptera). *Arquivo da Universidade Federal Rural do Rio de Janeiro* 7(2): 89–100.
- Piccinini, R.S. 1974. Lista provisória dos quirópteros da coleção do Museu Paraense Emílio Goeldi (Chiroptera). *Boletim do Museu Paraense Emílio Goeldi* 77(11): 1–32.
- Piccinini, R.S., A.L. Peracchi, S.D.L. Raimundo, A.M. Tannure, J.C.P. Souza, S.T. Albuquerque and L.L. Furtado. 1991. Observações sobre o hábito alimentar de *Diphylla ecaudata* Spix, 1923 (Chiroptera). *Revista Brasileira de Medicina Veterinária* 13(2): 8–10.
- Pinto-da-Rocha, R. 1995. Sinopse da fauna cavernícola do Brasil (1097–1994). *Papéis Avulsos de Zoologia* 39(6): 61–173.
- Pinto-da-Rocha, R., G. Sessego, and E. Sipinski. 2001. A fauna das Grutas de Botuverá, Santa Catarina, Brasil; pp. 135–153, in: L.F.S. Rocha, K.L. Oliveira and G.C. SESS (eds.). *Conservando Cavernas. Quinze anos de Espeleologia*. GEOP-Açungui. Curitiba: GEOP-AÇUNGUI.
- Pira, A. 1904. Über Fledermäuse von São Paulo. *Zoologische Anzeiger* 28(1): 12–19 (<http://biodiversitylibrary.org/page/9902709>).
- Reis, N.R., A.L. Peracchi, M.L. Sekiama, and I.P. Lima. 2000. Diversidade de morcegos (Chiroptera, Mammalia) em fragmentos florestais no estado do Paraná, Brasil. *Revista Brasileira de Zoologia* 17(3): 697–704 (doi: 10.1590/S0101-81752000000300015).
- Reis, N.R., M.L.S. Barbieri, I.P. Lima and A.L. Peracchi. 2003. O que é melhor para manter a riqueza de espécies de morcegos (Mammalia, Chiroptera), um fragmento florestal grande ou vários fragmentos de pequeno tamanho? *Revista Brasileira de Zoologia* 20(2): 225–230 (10.1590/S0101-81752003000200009).
- Reis, N.R., P.H. Gallo, A.L. Peracchi, I.P. Lima and M.N. Fregonezi. 2012. Sensitivity of populations of bats (Mammalia, Chiroptera) in relation to human development in northern Paraná, southern Brazil. *Brazilian Journal of Biology* 72(3): 511–518 (doi: 10.1590/S1519-69842012000300014).
- Ruschi, A. 1951. Morcegos do estado do Espírito Santo. Descrição de *Diphylla ecaudata* Spix e algumas observações a seu respeito. *Boletim do Museu de Biologia Prof. Mello Leitão* 6(3): 1–6.
- Ruschi, A. 1978. A atual fauna de mamíferos, aves e répteis da reserva biológica de Comboios. *Boletim do Museu de Biologia Prof. Mello Leitão* (90): 1–26.
- Ruschi, A. 1982. Mamíferos e aves observadas na Reserva Biológica de Pedra Azul no estado do Espírito Santo. *Boletim do Museu de Biologia Prof. Mello Leitão* (103): 1–15.
- Sanborn, C.C. 1949. Mammals from the Rio Ucayali, Peru. *Journal of Mammalogy* 30(3): 277–288.
- Sbragia, I.A. and A. Cardoso. 2008. Quiróptero-fauna (Mammalia: Chiroptera) cavernícola da Chapada Diamantina, Bahia, Brasil. *Chiroptera Neotropical* 14(1): 360–365.
- Sikes, R.S., W.L. Gannon, and the Animal Care and Use Committee of the American Society of Mammalogists. 2011. Guidelines of the American Society of Mammalogists for the Use of Wild Mammals in Research. *Journal of Mammalogy* 92(1): 235–253 (10.1644/10-MAMM-F-355.1).
- Solari, S., E. Vivar, P.M. Velazco, J.J. Rodríguez, D.E. Wilson, R.J. Baker, and J.L. Mena. 2001. The small mammal community of the Lower Urubamba Region, Peru; pp. 171–181, in: A. Alonso, F. Dallmeier, and P. Campbell (eds.). *Urubamba: The Biodiversity of a Peruvian Rainforest*. Washington: The Smithsonian Institution.
- Sousa, M., A. Langguth and E.A. Gimenez. 2004. Mamíferos dos brejos de altitude de Paraíba e Pernambuco; pp. 229–254, in: K.C. Porto, J.J.P. Cabral, and M. Tabarelli, (eds.). *Brejos de Altitude em Pernambuco e Paraíba: História Natural, Ecologia e Conservação*. Brasília: Ministério do Meio Ambiente.
- Stutz, W.H., M.C. Albuquerque, W. Uieda, E.M. Macedo and C.B. França. 2004. Updated list of bats from Uberlândia, state of Minas Gerais southeastern Brazil. *Chiroptera Neotropical* 10 (1–2): 188–90.
- Teixeira, S.C. and A.L. Peracchi. 1996. Morcegos do Parque Estadual da Serra da Tiririca, Rio de Janeiro, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 13(1): 61–66 (doi: 10.1590/S0101-81751996000100005).
- Thomas, O. 1900. List of the mammals obtained by Dr. G. Franco Grillo in the province of Parana, Brazil. *Annali del Museo Civico di Storia Naturale Genova, Série 2*, 20:546–49.
- Thomas, O. 1926. The Godman-Thomas Expedition to Peru III. On mammals collected by Mr. R. W. Hendee in the Chachapoyas region of north Peru. *Journal of Natural History, Series 9*, 18(103):156–167.
- Trajano, E. 1985. Ecologia de populações de morcegos cavernícolas em uma região cárstica do sudeste do Brasil. *Revista Brasileira de Zoologia* 2(5): 255–320 (doi: 10.1590/S0101-81751984000100001).
- Trajano, E. and E.A. Gimenez. 1998. Bat community in a cave from eastern Brazil, including a new record of *Lionycteris* (Phyllostomidae, Glossophaginae). *Studies on Neotropical Fauna & Environment* 33(2): 69–75 (doi: 10.1076/snfe.33.2.69.2156).
- Tuttle, M.D. 1970. Distribution and zoogeography of Peruvian bats, with comments on natural history. *The University of Kansas Science Bulletin* 49(2): 45–86.
- Uieda, W. 1986. Aspectos da morfologia lingual das três espécies de morcegos hematófagos (Chiroptera, Phyllostomidae). *Revista Brasileira de Biologia* 46(3): 581–87.
- Uieda, W. 1992. Período de atividade alimentar e tipos de presa dos morcegos hematófagos (Phyllostomidae) no sudeste do Brasil. *Revista Brasileira de Biologia* 52: 563–573.
- Wenzel, R.L., V.J. Tipton, and A. Kiewlicz. 1966. The streblid batflies of Panama (Diptera: Calyptridae: Streblidae); pp. 405–675, in: R.L. Wenzel and V.J. Tipton (eds.). *Ectoparasites of Panama*. Chicago: Field Museum of Natural History.

RECEIVED: December 2013

ACCEPTED: October 2014

PUBLISHED ONLINE: December 2014

EDITORIAL RESPONSIBILITY: Marcelo Nogueira